


NAVAL POSTGRADUATE SCHOOL

Monterey, California



A THEOREM ON GEOMETRIC PROBABILITY
AND APPLICATIONS

by

Peter C. C. Wang

20 November 1972

Approved for public release; distribution unlimited.

NAVAL POSTGRADUATE SCHOOL
Monterey, California

Rear Admiral Mason B. Freeman
Superintendent

M. U. Clauser
Provost

ABSTRACT:

Take any line ℓ in the plane and put a Poisson process of points with parameter λ on this line. At each point of the process construct an intersecting line, say ℓ_n at the n th point, by choosing the angle of intersection θ_n between ℓ and ℓ_n from an underlying distribution $F(\theta)$ such that the $\langle \theta_n \rangle$ are a sequence of independent and identically distributed random variables and additionally, independent of the Poisson process variables. Take any other line ℓ' , the intersections of ℓ' by the lines ℓ_n form a Poisson process with parameter

$$\lambda \left\{ \int_{-\infty}^0 F(\theta^*) ds + \int_0^{\infty} [1-F(\theta^*)] ds \right\}$$

where

$$\theta^* = \cos^{-1} [(-s + \cos \alpha)(1 + s^2 - 2s \cos \alpha)^{-1/2}]$$

and α is the angle between lines ℓ and ℓ' . The above result will imply both Rényi's and Breiman's results [1], [2]. A different but simplified proof of the Rényi result is also included.

This task was supported by: Office of Naval Research
Contract Number NR-042-286

NPS-53WG72111A

20 November 1972

A Theorem on Geometric Probability
and Applications

by

Peter C. C. Wang
Mathematics Department
Naval Postgraduate School
Monterey, California 93940

*The work was supported by the Office of Naval Research under Contract Number NR 042-286 and the Foundation Research at the Naval Postgraduate School.

A Theorem on Geometric Probability and Applications

ABSTRACT

Take any line ℓ in the plane and put a Poisson process of points with parameter λ on this line. At each point of the process construct an intersecting line, say ℓ_n at the n th point, by choosing the angle of intersection θ_n between ℓ and ℓ_n from an underlying distribution $F(\theta)$ such that the $\langle \theta_n \rangle$ are a sequence of independent and identically distributed random variables and additionally, independent of the Poisson process variables. Take any other line ℓ' , the intersections of ℓ' by the lines ℓ_n form a Poisson process with parameter

$$\lambda \left\{ \int_{-\infty}^0 F(\theta^*) ds + \int_0^{\infty} [1-F(\theta^*)] ds \right\}$$

where

$$\theta^* = \cos^{-1} [(-s + \cos \alpha)(1 + s^2 - 2s \cos \alpha)^{-1/2}]$$

and α is the angle between lines ℓ and ℓ' . The above result will imply both Rényi's and Breiman's results [1], [2]. A different but simplified proof of the Rényi result is also included.

A Theorem on Geometric Probability and Applications*

by

PETER C. C. WANG

NAVAL POSTGRADUATE SCHOOL, Monterey, California

1. Introduction

Rényi [1] has considered an interesting model of traffic flow on a divided highway which extends to infinity in one direction without traffic lights or other inhomogeneities. It is assumed that each car travels at a constant speed which is a random variable and passing is always possible without delays. Among others, Rényi has obtained some results regarding mainly the spatial distribution of cars along the highway when the temporal distribution of cars is assumed to be described by a Poisson Process. The purpose of this paper is to discuss a number of results that can be related to low density traffic flow models on an infinite highway. These models were initiated and developed principally in papers by Rényi [1], Weiss and Herman [2], Breiman [3] sometimes without specific reference to low density traffic flow, and these models are unified by Solomon and Wang [5] in a study of Non-homogeneous Poisson Fields of Random lines. Of the theorems presented in Section 2, all are known results [5] but the proofs are included for the sake of completeness. Main results are presented in Section 3. This main result will imply immediately the results reported in [5].

2. Low Density Traffic Flow Models

In this section, we shall provide a new proof of the Rényi theorems and include other results dealing with low density traffic flow. It will also be demonstrated for Rényi's model, that if the spatial distribution of cars is assumed to obey a Poisson process then the temporal distribution

*This work was supported by the Office of Naval Research under Contract Number NR 042-286 and the Foundation Research at the Naval Postgraduate School.

of cars (i.e., arrival times at some fixed position) is again a Poisson process. This result establishes a crucial structural property of Rényi's model for low-density traffic. Rényi found it convenient to start from the stochastic process of entrance times of the cars at a fixed point on the highway. Other authors start from the spatial process of cars distributed in locations along the infinite highway at some fixed time according to some random law. The Poisson process is the assumed machinery governing the car entrance times or equivalently car positions and the speed distributions for each car are assumed to be independently and identically distributed (i.i.d.) with a common distribution $G(V)$.

Starting from this spatial process, Breiman [3] considered the idealized model and proved that the Poisson process is the only process obeying the time-invariance property-namely, if at a time t_0 , the spatial process is Poisson with specific parameter, and the speeds of the cars are i.i.d. with respect to each other and the positions of the cars at time t_0 , then the process will have the same properties at any other time t . Similarly, we can obtain the spatial-invariance property-namely, if at a location x_0 , the temporal process is Poisson with specific parameter, and the speeds of the cars are i.i.d. with respect to each other and the entrance times of the cars at location x_0 , then the process will have the same properties at any other location x . These invariant properties insure that the randomness is unaffected by the choice of origin (i.e., randomness is unaffected by translation). Furthermore, the underlying process has independent stationary increments. It is obvious that the number of cars in a zero distance is zero and the number of cars arriving in a zero temporal interval is zero. In order to establish that the underlying process is Poisson, it is now sufficient to show that the number of cars in any temporal (or spatial)

interval has a Poisson distribution [7, p. 119].

In detail, the assumptions of Renyi's model are:

(i) Instants $\langle t_i \rangle_{i=1}^{\infty}$ at which cars enter the highway at a fixed position form a homogeneous Poisson process with parameter ω .

(ii) A car arriving at a certain point on the highway at instant t_i chooses a velocity V_i and then moves with this constant velocity. The random variables $\langle V_k \rangle$ are independently and identically distributed with distribution function $G(v) = \Pr\{V \leq v\}$ and sequences $\langle V_k \rangle$ and $\langle t_k \rangle$ are independent.

(iii) $\int_0^{\infty} \frac{1}{v} dG(v) < \infty$ i.e., the mean value of $\frac{1}{v}$ is finite; without this condition a traffic jam would arise and make all traffic flow impossible.

(iv) No delay in overtaking a car traveling at a slower speed when it is approached.

Suppose an arbitrary car $K(t_0, v_0)$ arrives at some fixed point of the highway at time t_0 where it assumes and maintains the fixed speed v_0 .

Let $\langle t_k^+ \rangle$ ($\langle t_k^- \rangle$) denote the instants at which the car $K(t_0, v_0)$ is overtaken by faster (overtaking slower) cars. Rényi has obtained the following results.

Theorem 1. (Rényi) The instants $\langle t_k^+ \rangle$ and $\langle t_k^- \rangle$ form two independent homogeneous Poisson processes, with parameters:

$$\omega^+(v_0) = \omega \int_0^{v_0} \frac{v_0 - v}{v} dG(v) \quad \text{and} \quad \omega^-(v_0) = \omega \int_{v_0}^{\infty} \frac{v - v_0}{v} dG(v).$$

Rényi's proof of the above theorem is based on the following two properties of Poisson processes:

(A) If $\langle t_i \rangle$ are the instants of time when an event occurs in a homogeneous Poisson process with parameter ω , and ζ_1, ζ_2, \dots is a sequence

of independent positive random variables, each having the same distribution function $G(\zeta)$ and each independent of the process $\langle t_k \rangle$, then the time instants $t_k \zeta_k$ ($k = 1, 2, \dots$) also form a homogeneous Poisson process with density

$$\omega^* = \omega \int_0^\infty \frac{1}{\zeta} dG(\zeta).$$

(B) If a subsequence $\langle t_{v_k} \rangle$ of the instants $\langle t_k \rangle$, in which an event occurs in a Poisson process with density ω , is selected at random in such a way that for each j the probability of the event A_j that j should belong to the subsequence $\langle v_k \rangle$ is equal to r ($0 < r < 1$) and the events A_j ($j = 1, 2, \dots$) are independent; and if $\langle t_{u_k} \rangle$ are the instants that are not selected, (i.e., j belongs to the sequence $\langle u_k \rangle$ if and only if it does not belong to the sequence $\langle v_k \rangle$), then $\langle t_{v_j} \rangle$ and $\langle t_{u_k} \rangle$ are two independent Poisson processes with density ωr and $\omega(1-r)$.

It is now known from a result of Wang [6] that property (B) is a characteristic property for Poisson processes. We shall establish Theorem 1 without using property (A). It can be shown that property B implies property A (see Theorem 2 below).

Proof of Theorem 1. The trajectory of any car in time-space (Diagram 1)


for Renyi's low density traffic model is realized by a straight line.

Let us denote the trajectories of all cars on the highway as a set A .

Denote M_L^+ the number of lines in A that intersect segment L of the trajectory of car $K(0, v_0)$ from below and M_p the number of lines in A whose arrival times are in $(0, t_0)$. Then

Diagram 1.

Time-Space Diagram



$$\begin{aligned}
\Pr\{M_L^+ = n\} &= \sum_{m=n}^{\infty} \binom{m}{n} \mu^n (1-\mu)^{m-n} e^{-\omega t_0} \frac{(\omega t_0)^m}{m!} \\
&= e^{-\omega t_0} \mu \frac{(\omega t_0 \mu)^n}{n!}
\end{aligned}$$

where

$$\mu = \Pr\{M_L^+ = 1 \mid M_p = 1\}$$

$$= \frac{1}{t_0} \int_0^{t_0} \Pr\{V \geq \frac{x_0}{t_0 - p}\} dp$$

$$= \int_{v_0}^{\infty} \frac{v - v_0}{v} dG(v), \text{ where } v_0 = \frac{x_0}{t_0}.$$

Similarly, we define M_L^- as the number of lines in A intersecting L from above and M_p^c as the number of lines in A whose arrival times fall in the interval $-c(c > 0)$ and 0. We can compute

$$\Pr\{M_L^- = n\} = \lim_{c \rightarrow \infty} e^{-\omega c \mu_c^*} \frac{(\omega c \mu_c^*)^n}{n!}$$

where

$$\mu_c^* = \Pr\{M_L^- = 1 \mid M_p^c = 1\}.$$

$$\lim_{c \rightarrow \infty} c \mu_c^* = \lim_{c \rightarrow \infty} c \Pr\{M_L^- = 1 \mid M_p^c = 1\}$$

$$= \lim_{c \rightarrow \infty} \int_{-c}^0 \int_0^{\frac{x_0}{t_0 - p}} dG(v) dp$$

$$= \int_{-\infty}^0 \int_0^{\frac{x_0}{t_0 - p}} dG(v) dp$$

$$= \int_0^{v_0} \frac{x_0 - vt_0}{v} dG(v)$$

$$= t_0 \int_0^{v_0} \frac{v_0 - v}{v} dG(v) .$$

Random variables M_L^- and M_L^+ are independent because the events involved come from disjoint intervals. This completes the proof of Theorem 1.

The counting interval employed in the above theorem is on the time axis.

In what follows, a similar approach to the problem dealing with a spatial counting interval is employed and produces some interesting results.

Denote M^- the number of lines in A intersecting L from above and

$M_{x_0}^c$ for the number of lines in A whose spatial positions at $t = 0$

are between 0 and $-c$. Let λ^* be the spatial density.

Then we have

$$\Pr\{M^- = n\} = e^{-\lambda^* x_0 \mu} \frac{(\lambda^* x_0 \mu)^n}{n!}$$

where

$$\mu = \Pr\{M^- = 1 \mid M_{x_0}^c = 1\}$$

$$= \frac{t_0}{x_0} \int_0^{v_0} (v_0 - v) dG(v) ;$$

and

$$\begin{aligned} \Pr\{M^+ = n\} &= \lim_{c \rightarrow \infty} \sum_{m=n}^{\infty} \Pr\{M^+ = n \mid M_{x_0}^c = m\} \Pr\{M_{x_0}^c = m\} \\ &= \lim_{c \rightarrow \infty} e^{-\lambda * c \mu_1^c} \frac{(\lambda * c \mu_1^c)^n}{n!} \end{aligned}$$

where

$$\mu_1^c = \Pr\{M^+ = 1 \mid M_{x_0}^c = 1\} .$$

It can be easily verified that

$$\lim_{c \rightarrow \infty} c \mu_1^c = t_0 \int_{v_0}^{\infty} (v - v_0) dG(v) ,$$

and random variables M^- and M^+ are independent. Now denote $M = M^+ + M^-$ and we conclude that

$$\Pr\{M = n\} = e^{-\lambda * t_0 \int_0^{\infty} |v_0 - v| dG(v)} \cdot \frac{[\lambda * t_0 \int_0^{\infty} |v_0 - v| dG(v)]^n}{n!} .$$

The above result appeared initially in a paper by Weiss and Herman[2] who arrived at it from different considerations.

In the next paragraph, results are stated about the spatial distribution of vehicles if the temporal process is assumed to be Poisson.

Denote S^+ the number of lines in A intersecting $(0, x_0)$ and $x_0 > 0$ at time zero and S_c^+ the number of lines in A whose arrival times are in the interval $(-c, 0)$. We further denote S^- the number of lines in A intersec-

ting $(-x_0, 0)$, $x_0 > 0$ at time zero and S_c^- the number of lines in A whose arrival times are in the interval $(0, c)$. Let us compute quantities: $\Pr\{S^+ = n\}$, $\Pr\{S^- = n\}$ and $\Pr\{S = S^+ S^- = n\}$.

$$\Pr\{S^+ = n\} = \lim_{c \rightarrow \infty} e^{-\omega c \mu_2} \frac{(\omega c \mu_2)^n}{n!}$$

where ω is the temporal density and

$$\mu_2 = \Pr\{S^+ = 1 \mid S_c^+ = 1\}.$$

It can be shown easily that

$$\lim_{c \rightarrow \infty} c \mu_2 = x_0 \int_0^\infty \frac{1}{v} dG(v).$$

We conclude that

$$\Pr\{S^+ = n\} = e^{-\omega x_0 \int_0^\infty \frac{1}{v} dG(v)} \frac{[\omega x_0 \int_0^\infty \frac{1}{v} dG(v)]^n}{n!}.$$

Similarly, we obtain

$$\Pr\{S^- = n\} = \Pr\{S^+ = n\},$$

and

$$\Pr\{S = S^+ + S^- = n\} = e^{-2\omega x_0 \int_0^\infty \frac{1}{v} dG(v)} \frac{[2\omega x_0 \int_0^\infty \frac{1}{v} dG(v)]^n}{n!}.$$

We can now summarize as follows.

Theorem 2. If $\langle t_i \rangle$ forms a Poisson process with parameter ω and sequences $\langle t_i \rangle$ and $\langle V_i \rangle$ are independent then the locations of vehicles on the highway at time $t = 0$ namely $\langle x_i \rangle$ forms a Poisson process with parameter

$$\omega \int_0^\infty \frac{1}{v} dG(v)$$

Theorem 3. If $\langle x_i \rangle$ forms a Poisson process with parameter λ^* and sequences $\langle x_i \rangle$ and $\langle V_i \rangle$ are independent and $\langle x_i^+ \rangle$ denotes the positions at which the car $K(0, v_0)$ overtakes slower cars and $\langle x_i^- \rangle$ denotes the positions at which car $K(0, v_0)$ is overtaken by faster cars; then the two sequences $\langle x_i^+ \rangle$ and $\langle x_i^- \rangle$ form two independent (homogeneous) Poisson processes, with parameters:

$$\lambda_+^*(v_0) = \lambda^* \int_0^{v_0} (v_0 - v) dG(v) \quad \text{and} \quad \lambda_-^*(v_0) = \lambda^* \int_{v_0}^{\infty} (v - v_0) dG(v) \quad .$$

This result is analogous to the Rényi result which we developed as Theorem 1 except that the counting of overtakings is accomplished on the spatial axis rather than on the time axis. The next theorem provides a result analogous to that in Theorem 2.

Theorem 4. If $\langle x_i \rangle$ forms a Poisson process with parameter λ^* and sequences $\langle x_i \rangle$ and $\langle V_i \rangle$ are independent and $\langle V_i \rangle$ are i.i.d. random variables with common distribution $G(v) = \Pr \{V \leq v\}$; then the corresponding $\langle t_i \rangle$ arrival times at position $x = 0$ forms a Poisson process with parameter $\lambda^*E(V)$ where

$$E(V) = \int_0^{\infty} v dG(v).$$

Proof. The proof is again based on the binomial mixing as presented in Property (B) and hence details are omitted.

3. Main Results

It is the purpose of this section to establish a result (Theorem 5 below) that by applying property (B) to it would imply both Rényi's and Breiman's results.

Let us take any line ℓ in the plane and put a Poisson process of points

on this line. At each point of the process construct an intersecting line ℓ_n at the n -th point S_n by choosing the angle of intersection θ_n between lines ℓ and ℓ_n from an underlying distribution $F(\theta)$ such that the $\langle \theta_n \rangle$ are a sequence of independent and identically distributed random variables and additionally, independent of the Poisson process. Now, take any other line ℓ' and look at the intersections of line ℓ' by the lines ℓ_n . We wish to show that the intersection points $\langle S'_n \rangle$ form a Poisson process and the sequence $\langle \theta'_n \rangle$ of the consecutive angles of intersection is independent and identically distributed. It is clear that if the line ℓ' is parallel to the line ℓ , then the intersection points $\langle S'_n \rangle$ of lines ℓ_n and ℓ' is again Poisson with the same parameter as the process on the line ℓ because of the invariance property mentioned in section 1. We can now state the following.

Theorem 5. If $\langle S_n \rangle$ forms a Poisson process with parameter λ on line ℓ and sequences $\langle \theta_n \rangle$ are i.i.d. with a common distribution $F(\theta)$, $0 \leq \theta < \pi$, and furthermore, the sequences $\langle S_n \rangle$ and $\langle \theta_n \rangle$ are independent, then

(i) the sequence $\langle S'_n \rangle$ forms a Poisson process on line ℓ' with parameter

$$\lambda^* = \lambda \left\{ \int_{-\infty}^0 F(\theta^*) ds_n + \int_0^{\infty} [1 - F(\theta^*)] ds_n \right\}$$

where

$$\theta^* = \cos^{-1} [(-s_n + \cos \alpha) (1 + s_n^2 - 2s_n \cos \alpha)^{-1/2}]$$

and α is the angle between lines ℓ and ℓ' , and

(ii) the angles $\langle \theta'_n \rangle$ formed between lines ℓ' and ℓ_n are i.i.d. with common distribution $F(\theta' + \alpha)$, $-\alpha \leq \theta' < \pi - \alpha$.

Since the counting measure on line ℓ is invariant under translation and the

corresponding process has independent stationary increments, it is sufficient to show that the sequences $\langle S'_n \rangle$ forms a Poisson process by showing that the counting measure has a Poisson distribution. i.e., to show that

$$\Pr\{N_{\ell'} = k\} = e^{-\lambda^*} \frac{(\lambda^*)^k}{k!}; \quad k = 0, 1, 2, \dots$$

where the random variable $N_{\ell'}$ denotes the number of intersections between lines ℓ_n and ℓ' in a given unit interval on line ℓ' .

Proof of Theorem 5.

Because of the invariance property mentioned in section one, it is sufficient to choose the intersection of lines ℓ' and ℓ as the origin and denote the random variable N_{ℓ}^c the number of points in the interval $(-c, c)$ on line ℓ . Then we have

$$\begin{aligned} \Pr\{N_{\ell'} = k\} &= \lim_{c \rightarrow \infty} \sum_{m=k}^{\infty} \Pr\{N_{\ell'} = k | N_{\ell}^c = m\} \Pr\{N_{\ell}^c = m\} \\ &= \lim_{c \rightarrow \infty} e^{-2c\lambda\mu} \frac{(2c\lambda\mu)^k}{k!}; \quad k = 0, 1, 2, \dots, \end{aligned}$$

where

$$\mu = \Pr\{N_{\ell'} = 1 | N_{\ell}^c = 1\}.$$

Upon evaluation of μ we have

$$\lim_{c \rightarrow \infty} 2c\mu = \int_{-\infty}^0 F(\theta^*) dS_n + \int_0^{\infty} [1-F(\theta^*)] dS_n.$$

Furthermore, the angles $\langle \theta'_n \rangle = \langle \theta_n - \alpha \rangle$ (see diagram 2) are clearly i.i.d. with common distribution $F(\theta' + \alpha)$ for $-\alpha \leq \theta < \pi - \alpha$. This completes the proof of the theorem.




Diagram 2

It is reasonable to conjecture that the sequences $\langle \theta'_n \rangle$ and $\langle S'_n \rangle$ formed in Theorem 5 are independent sequences, however it is not true in general unless the lines $\langle \ell_n \rangle$ are all parallel lines. To see this, we can use the following indirect argument:

Based on the results stated in Theorem 2 and Theorem 4 plus the independence condition needed, one might expect to get the following identity

$$\omega = \omega \int_0^\infty \frac{1}{v} dG(v) \int_0^\infty v dG(v)$$

But the identity is true if and only if all cars are travelling at the same speed.

Applying property B to Theorem 5 and restricting $\alpha = \pi/2$ and $0 \leq \theta < \frac{\pi}{2}$ will result Rényi's Theorem. Other theorems stated in section 2 are implied by Theorem 5 in a similar manner.

References:

1. Rényi, A. - On Two Mathematical Models of the Traffic On A Divided Highway. J. Appl. Prob. 1, (1964) 311-320
2. Weiss, George and Herman, Robert - Statistical Porperties of Low-Density Traffic. Quarterly of Applied Mathematics XX, (1962) 121-130
3. Breiman, L. - The Poisson Tendency In Traffic Distribution. Ann. Math. Stat. 34, (1963) 308-311
4. Wang, Peter C. -C. - On A Theorem of Rényi and Extensions. Notices of Amer. Math. Soc. 17 (1970) 640
5. Solomon, H. and Wang, Peter C. -C. - Non-homogeneous Poisson Fields of Random Lines With Applications. Sixth Berkeley Symposium on Math. Stat. and Probability. University of California, Berkeley (to appear)
6. Wang, Peter C. -C. - A Characterization of the Poisson Distribution Based on Random Splitting and Random Expanding. Technical Report No. 160, Dept. of Statistics, Stanford University. August, 1970
7. Parzen, Emanuel - Stochastic Processes Holden-Day, 1965
8. Breiman, L. - On Some Probability Distributions Occurring in Traffic flow, Bull. Int. Statistical Institute, 39 (1962) 155-161

DISTRIBUTION LIST

	Copies		Copies
Statistics and Probability program		Office of Naval Research Branch Office	
Office of Naval Research		536 South Clark Street	
Attn: Dr. B. J. McDonald		Attn: Dr. Robert Buchal	
Arlington, Virginia 22217	3	Chicago, Illinois 60605	1
Director, Naval Research Laboratory		Director	
Attn: Library, Code 2029		Office of Naval Research	
(ONRL)		Branch Office	
Washington, D. C. 20390	6	1030 East Green Street	
		Attn: Dr. A. R. Laufer	
		Pasadena, California 91101	1
Defense Documentation Center		Office of Naval Research	
Cameron Station		Branch Office	
Alexandria, Virginia 22314	12	1030 East Green Street	
Defense Logistics Studies		Attn: Dr. Richard Lau	
Information Exchange		Pasadena, California 91101	1
Army Logistics Management Center		Office of Naval Research	
Attn: Arnold Hixon		San Francisco Area Office	
Fort Lee, Virginia 23801	1	50 Fell Street	
Technical Information Division		San Francisco, California 94102	1
Naval Research Laboratory		Technical Library	
Washington, D. C. 20390	6	Naval Ordnance Station	
Office of Naval Research		Indian Head, Maryland 20640	1
New York Area Office		Naval Ship Engineering Center	
207 West 24th Street		Philadelphia	
Attn: Dr. Jack Laderman		Division Technical Library	
New York, New York	1	Philadelphia, Pennsylvania 19112	1
Director		Bureau of Naval Personnel	
Office of Naval Research		Department of the Navy	
Branch Office		Technical Library	
495 Summer Street		Washington, D. C. 20370	1
Attn: Dr. A. L. Powell		Library, Code 0212	
Boston, Massachusetts 02210	1	Naval Postgraduate School	
Director		Monterey, California 93940	1
Office of Naval Research		Library	
Branch Office		Naval Electronics Laboratory Center	
536 South Clark Street		San Diego, California 92152	1
Attn: Dr. A. R. Dawe			
Chicago, Illinois 60605	1		

	Copies		Copies
Naval Undersea Center Technical Library San Diego, California 92132	1	Princeton University Department of Statistics Attn: Prof. G. S. Watson Princeton, New Jersey 05840	1
Applied Mathematics Laboratory Naval Ships Research and Development Center Attn: Mr. Gene H. Gleissner Washington, D. C. 20007	1	Stanford University Department of Statistics Attn: Prof. T. W. Anderson Stanford, California 94305	1
Office of Chief of Naval Operations (Op 964) Ballston Tower No. 2 Attn: Mr. A. S. Rhodes Arlington, Virginia 22203	1	University of California Department of Statistics Attn: Prof. P. J. Bickel Berkeley, California 94720	1
Naval Ships Systems Command Ships 0311 National Center No. 3 Attn: Miss B. S. Orleans Arlington, Virginia 20360	1	University of Washington Department of Mathematics Attn: Prof. Z. W. Birnbaum Seattle, Washington 98105	1
University of Chicago Department of Statistics Attn: Prof. W. Kruskal Chicago, Illinois 60637	1	Harvard University Department of Statistics Attn: Prof. W. G. Cochran Cambridge, Massachusetts 02139	1
Stanford University Department of Operations Research Attn: Prof. G. J. Lieberman Stanford, California 94305	1	Columbia University Department of Civil Engineering and Engineering Mechanics Attn: Prof. C. Derman New York, New York 10027	1
Florida State University Department of Statistics Attn: Prof. I. R. Savage Tallahassee, Florida 32306	1	Columbia University Department of Mathematics Attn: Prof. H. Robins New York, New York 10027	1
Florida State University Department of Statistics Attn: Prof. R. A. Bradley Tallahassee, Florida 32306	1	New York University Institute of Mathematical Science Attn: W. M. Hirsch New York, New York 10453	1
Princeton University Department of Statistics Attn: Prof. J. W. Tukey Princeton, New Jersey 08540	1	University of North Carolina Department of Statistics Attn: Prof. W. L. Smith Chapel Hill, North Carolina 27514	1
		Dean of Research Naval Postgraduate School Monterey, California 93940	2

	Copies		Copies
Clemson University Department of Mathematics Attn: Prof. K. T. Wallenius Clemson, South Carolina 29631	1	Program in Logistics The George Washington University Attn: Dr. W. H. Marlow 707 22nd Street, N. W. Washington, D. C. 20037	1
University of California Department of Statistics Attn: Charles E. Antoniak Berkeley, California 94720	1	Mississippi Test Facility Earth Resources Laboratory (Code GA) Attn: Mr. Sidney L. Whitley Bay St. Louis, Mississippi 39520	1
Clarkson College of Technology Division of Research Attn: Prof. M. Arozullah Potsdam, New York 13676	1	Naval Postgraduate School Department of Operations Research and Administrative Sciences Attn: Prof. P. A. W. Lewis Monterey, California 93940	1
University of Southern California Electrical Sciences Division Attn: Prof. W. C. Lindsey Los Angeles, California 90007	1	Southern Methodist University Department of Statistics Attn: Prof. W. R. Schucany Dallas, Texas 75222	1
Case Western Reserve University Department of Mathematics and Statistics Attn: Prof. S. Zacks Cleveland, Ohio 44106	1	Webb Institute of Naval Architecture Attn: Prof. O. J. Karst Crescent Beach Road Glen Cove, New York 11543	1
University of Florida Department of Electrical Engineering Attn: Prof. D. G. Childers Gainesville, Florida 32601	1	University of Missouri Department of Statistics Attn: Prof. W. A. Thompson, Jr. Columbia, Missouri 65201	1
Stanford University Department of Statistics Attn: Prof. H. Chernoff Stanford, California 94305	1	Rice University Department of Mathematical Sciences Attn: Prof. J. R. Thompson Houston, Texas 77001	1
Naval Research Laboratory Electronics Division (Code 5267) Attn: Mr. Walton Bishop Washington, D. C. 20390	1	University of California System Science Department Attn: Prof K. Yao Los Angeles, California 90024	1
Commandant of the Marine Corps (Code AX) Attn: Dr. A. L. Slafkosky Scientific Advisor Washington, D. C. 20380	1	Naval Postgraduate School Department of Mathematics Attn: P. C. C. Wang Monterey, California 93940	20

	Copies		Copies
Raytheon Company Submarine Signal Division Attn: Dr. W. S. Liggett, Jr. Portsmouth, Rhode Island 02971	1	Smithsonian Institution Astrophysical Observatory Attn: Dr. C. A. Lundquist Cambridge, Massachusetts 02138	1
Systems Control, Inc. Attn: Dr. L. P. Seidman 260 Sheridan Avenue Palo Alto, California 44306	1	Naval Postgraduate School Department of Operations Research and Administrative Sciences Attn: Prof. J. D. Esary Monterey, California 93940	1
University of California Department of Information and Computer Science Attn: Prof. E. Masry La Jolla, California 92037	1	Polytechnic Institute of Brooklyn Department of Electrical Engineering Attn: Prof. M. L. Shooman Brooklyn, New York 11201	1
University of California School of Engineering Attn: Prof. N. J. Bershad Irvine, California 92664	1	Union College Institute of Industrial Administration Attn: Prof. L. A. Aroian Schenectady, New York 12308	1
University of California School of Engineering and Applied Science Attn: Prof. I. Rubin Los Angeles, California 90024	1	Ultrasystems, Inc. Attn: Dr. D. C. Dorrough 500 Newport Center Drive Newport Beach, California 92660	1
Virginia Polytechnic Institute Department of Statistics Attn: Prof. C. Kramer Blacksburg, Virginia 24061	1	University of New Mexico Department of Mathematics and Statistics Attn: Prof. W. J. Zimmer Albuquerque, New Mexico 87106	1
New York University Department of Electrical Engineering Attn: Prof. I. Yagoda Bronx, New York 10453	1	Princeton University Department of Statistics Attn: Prof. G. Simon Princeton, New Jersey 08540	1
University of Rochester Department of Statistics Attn: Prof. J. Keilson Rochester, New York 14627	1	Naval Ordnance Systems Command, NORD 035 Attn: Mr. O. Seidman Room 6E08, National Center #2 Arlington, Virginia 20360	1
University of Michigan Department of Industrial Engineering Attn: Prof. R. L. Disney Ann Arbor, Michigan 48104	1	Naval Coastal Systems Laboratory Code P761 Attn: Mr. C. M. Bennett Panama City, Florida 32401	1
Cornell University Department of Computer Science Attn: Prof. J. E. Hopcroft Ithaca, New York 14850	1	Food and Drug Administration Statistics and Information Science Division Health Protection Branch Attn: Dr. A. Petrasovits, Head, Survey Design and Quality Control 355 River Road, 4th Floor Vanier, Ontario, Canada	1

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author)

Naval Postgraduate School
Monterey, California 93940

2a. REPORT SECURITY CLASSIFICATION

UNCLASSIFIED

2b. GROUP

3. REPORT TITLE

A Theorem on Geometric Probability and Applications

4. DESCRIPTIVE NOTES (Type of report and, inclusive dates)

Scientific

5. AUTHOR(S) (First name, middle initial, last name)

Peter C. C. Wang

6. REPORT DATE

20 November 1972

7a. TOTAL NO. OF PAGES

24

7b. NO. OF REFS

8

8a. CONTRACT OR GRANT NO.

NR-042-286

b. PROJECT NO.

9a. ORIGINATOR'S REPORT NUMBER(S)

NPS-53WG72111A

c.

9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)

d.

10. DISTRIBUTION STATEMENT

Approved for public release; distribution unlimited.

11. SUPPLEMENTARY NOTES

12. SPONSORING MILITARY ACTIVITY

Office of Naval Research

13. ABSTRACT

Take any line ℓ in the plane and put a Poisson process of points with parameter λ on this line. At each point of the process construct an intersecting line, say ℓ_n at the n th point, by choosing the angle of intersection θ_n between ℓ and ℓ_n from an underlying distribution $F(\theta)$ such that the $\langle \theta_n \rangle$ are a sequence of independent and identically distributed random variables and additionally, independent of the Poisson process variables. Take any other line ℓ' , the intersections of ℓ' by the lines ℓ_n form a Poisson process with parameter

$$\lambda \left\{ \int_{-\infty}^0 F(\theta^*) ds + \int_0^{\infty} [1-F(\theta^*)] ds \right\}$$

where

$$\theta^* = \cos^{-1} [(-s + \cos \alpha)(1 + s^2 - 2s \cos \alpha)^{-\frac{1}{2}}]$$

and α is the angle between lines ℓ and ℓ' . The above result will imply both Rényi's and Breiman's results [1], [2]. A different but simplified proof of the Rényi result is also included.

14

KEY WORDS

LINK A

LINK B

LINK C

ROLE

WT

ROLE

WT

ROLE

WT

Geometric Probability

Poisson Process

Road Traffic Flow

Random lines

U150014

DUDLEY KNOX LIBRARY - RESEARCH REPORTS



5 6853 01057949 3

U1500